

ABORTION HEALTH PROFILE SIERRA LEONE 2019

| 1. General Statistics | | | 2. Legal status of abortion | | | |
|--|--------------------------------------|---|--|-------------------------|------------|--|
| Total population (000's), 2019 | | 7,813 | On request | Not specif | Yes | |
| · · ·································· | | 7,010 | On request For economic or social reasons | 0 | | |
| Total female population (15 - 49 years) (000's), 2019 Adolescent birth rate (per 1000 women aged 15-19 years), 2016 | | 1,915 | In cases of foetal impairment | 0 | | |
| | | | In cases of incest | 0 | | |
| | | 101 | In cases of intellectual/cognitive disability of the woman | 0 | | |
| Unintended pregnancies (000's), 2019 | | 205 | In cases of rape | 0 | | |
| | | 200 | To preserve a woman's health | 0 | | |
| Proportion of women who make their own informed decisions regarding sexual relations (% of women aged 15-49 years), 2013 | | 79 | To preserve a woman's mental health | 0 | | |
| Total fertility rate 15-49 (births per woman), 2016 | | | To preserve a woman's physical health To save a woman's life | 0 | | |
| | | 4 | Other | 0 | ~ | |
| 3. Causes of maternal de | oth (2017) | | | 10 2017 | <u>\</u> | |
| | alli (2017) | | 4. Maternal mortality ratio (20 1,400 1,360 | 10-2017 |) | |
| Maternal deaths aggravated by HIV/AIDS 0.8% | | | 1 190 | 1,180 | | |
| Late maternal deaths 1.0% | | | 1,200 1,270 1,210 1,180 | 1,120 |) 1,1 | |
| Maternal obstructed labor and uterine rupture 5.1 | % | | (1,200 1,210 1,210 1,180 1,210 1,180 0 1,180 0 0 1,000 1,00 | 1,120 | , 1, | |
| Maternal abortion and miscarriage 6. | 2% | | 5 800 | | | |
| | 4% | | 600 | | | |
| 1 1 3 7 | | | ¥ 400 | | | |
| Other maternal disorders | 7.4% | | | | | |
| Maternal hypertensive disorders 7.4% | | | 200 SDG country target 2030 = 140 | | | |
| Maternal sepsis and other maternal infections 18.7% | | | 2010 2011 2012 2013 2014 | 2015 2016 | 5 201 | |
| Maternal hemorrhage 19.54 | | | 5. Annual average rate of reduction: | | | |
| Indirect maternal deaths | | 27.6% | 27.6% 2010-2017 vs 2010-2030 | | | |
| 0% 5% 10% 15% 20 | | | Current AARR in MMR 2.8% | | | |
| 078 378 | % of Tota | | Expected AARR to reach MMR target 2030 | | 5.59 | |
| 6. FP indicators and unsafe ab | ortions | 7 Δνα | ilable in National Essential 8. | Post-abo | rtion | |
| averted due to modern contraceptive use | | | Medicines List - (2010) contraceptive | | | |
| averted due to modern contraceptive dse | | | | acceptan | | |
| | 69 | ³ Misoprostol | × – | | | |
| 60 60 64 | 03 | Mifemieteme | ~ | | | |
| $\begin{array}{cccccccccccccccccccccccccccccccccccc$ | | Mifepristone | × | No data | | |
| | | Misoprostol + Mifepristone combination | | | | |
| $\overline{38}$ 27 38 27 27 27 27 27 27 27 27 | 27 | 8 | References | | | |
| 20 18 20 22 24 25 26 | 27 27 | The formation of the second seco | | | | |
| 20 | | data acquired via web | site in July 2020; DHS and national surveys compiled in WHO Global Observat ; ICF, 2015. The DHS Program STATcompiler. Funded by USAID. Available fro | ory and UNFPA glo | bal databa | |
| 0 | | Accessed July 2020 | | m. <u>mp.//www.stab</u> | compiler.c | |
| 2012 2013 2014 2015 2016 2017 | 2018 2019 | 3: Global Burden of Di | cy Database (<u>https://abortion-policies.srhr.org/)</u> accessed in July 2020 sease Study 2017 (GBD 2017) Results.Seattle, United States: Institute for Hea | Ith Metrics and Eva | luation | |
| Year | | | Available from http://ghdx.healithdata.org/gbd-results-tool 4: Trends in maternal mortality 2000 to 2017: estimates by WHO, UNICEF, UNFPA, World Bank Group and the United Nations | | | |
| _egend ■ Contraceptive prevalence - modern methods (%) | | Population Division | | | | |
| Demand satisfied for modern contraception (%) Unmet need for family planning (%) | | | From FP2020 2019 Progress Report, most recent national surveys (DHS,PMA2020, MICS,RHS and other national surveys). Att.<u>http://progress.familyplanning2020.org/resources</u> | | | |
| Unsafe abortions averted (000's) | | 7: Global Abortion Pol | cy Database (https://abortion-policies.srhr.org/) accessed in July 2020 | | | |
| | | | Definitions and abbreviations | | | |
| Poproductive Maternal Health & Agoing (PMH) reference period as | | | averted due to modern contraceptive use: Number of unsafe abortions that did not occur during a specifie | | | |
| | | a result of the protection provided by modern contraceptive use during the reference period. valence - Modern Methods: Percentage of women of reproductive age who are using (or whose partner is | | | | |
| | Contracentive n | evalence Modern | ntraceptive method at a particular point in time | | | |
| Programme (afrgofrhrwh@who.int), Universal Health | using) a modern o | contraceptive method | at a particular point in time | • | | |
| | using) a modern o Unmet need: per | contraceptive method centage of fecund wo | | ne having the n | | |